

## Contents

<b>1</b>	<b>A Backward Monte Carlo algorithm to solve the Vector Radiative Transfer Equation</b>	<b>13</b>
1.1	The concept of Backward Monte Carlo	13
1.2	Performance of the BMC algorithm	15
1.3	Applications of the BMC algorithm	16
<b>2</b>	<b>Characterization of close-in exoplanet atmospheres with optical phase curves</b>	<b>37</b>
2.1	Kepler-7b's phase curve	38
2.2	The toy model for interpretation	38
2.3	Findings and perspectives	39
<b>3</b>	<b>High-dispersion spectro-polarimetry. A new approach for exoplanet characterization</b>	<b>67</b>
3.1	HDSP explained	68
3.2	Findings and perspectives	70
<b>4</b>	<b>Forward scattering from Titan's atmosphere</b>	<b>83</b>
4.1	Titan's phase curves from Cassini observations	83
4.2	Radiative transfer modeling	84
4.3	Other findings. Perspectives	85
<b>5</b>	<b>Forward scattering from exoplanet atmospheres</b>	<b>93</b>
5.1	An analytical theory for forward scattering from planetary atmospheres	93
5.2	Detectability of the brightness rise at exoplanets	94
<b>6</b>	<b>Examples of application of the BMC algorithm</b>	<b>115</b>