

Contents

1	Nonlinear propagation and absorption in wide band gap dielectrics	1
1.1	From linear to nonlinear propagation in a dielectric medium	1
1.2	Dispersion of a ultrashort laser pulse	5
1.3	Pulse propagation effects of the Kerr nonlinearity	6
2	Strong electric field excitation of wide band gap material	11
2.1	Primary absorption mechanisms	11
2.2	Secondary excitation and relaxation of the electron gas.	21
2.3	Transient optical response upon strong field ionization	24
3	Experimental	31
3.1	Glass principles and manufacturing	31
3.2	Laser pulse properties	39
3.3	The irradiation setup	42
4	Ablation threshold	47
4.1	Motivation	47
4.2	The ablation threshold fluence corresponds to kind and concentration of the network modifying ions	50
4.3	The ablations depth	52
4.4	The crater topography	56
4.5	Material specific ablation criterion: the dissociation energy	59
5	Nonlinear absorption and refraction	63
5.1	Z-scan experimental	63
5.2	Empirical calculations of the nonlinear refractive index	66
5.3	Measurements of the nonlinear refraction from z-scan	68
5.4	Measurement of the nonlinear absorption from z-scan	70
5.5	Transient refractive index changes measured by self-reflectivity	73
6	X-ray absorption spectroscopy of the structural changes induced by fs- laser irradiation	77
6.1	Principles of XANES	77
6.2	The PTB beamline and the chamber	78
6.3	Sample preparation	79
6.4	Setup geometry	81
6.5	Fs-laser induced ablation leads to a shift in the K-absorption edge . .	82

6.6	Discussion	84
6.7	The debris of a potassium-magnesium silicate glass generated by fs-laser induced ablation	87
7	Raman spectroscopic investigations	93
7.1	Raman spectroscopy experimental	94
7.2	Raman spectra of the non-irradiated glasses	95
7.3	Structural changes of the residual surface material after fs-laser ablation	100
8	Acknowledgements	105
	Bibliography	107