

Table of Contents

Abstract	II
Zusammenfassung	III
Abbreviations.....	V
1. Introduction	1
1.1 The bone marrow.....	1
1.1.1 Bone development	1
1.1.2 Bone and bone marrow structure.....	2
1.1.3 Bone cells	4
1.2 Hematopoietic stem cells.....	5
1.2.1 Origin and development	6
1.2.2 Hematopoietic stem cell niche	7
1.2.3 Hematopoiesis	11
1.3 Alternatives to animal testing.....	14
1.3.1 In vitro two-dimensional HSC culture	15
1.3.2 Tissue Engineering	17
1.3.3 Bone marrow tissue engineering	18
1.3.4 Organs-on-a-chip.....	20
2. Aim.....	23
3. Materials and Methods	24
3.1 Materials	24
3.1.1 Cell sources	24
3.1.2 Cell culture media and supplements.....	24
3.1.3 Buffers and reagents	25
3.1.4 Consumables	25
3.1.5 Antibodies for flow cytometry and immunofluorescence	26
3.1.6 Kits	28
3.1.7 qPCR Primers	28
3.1.8 Multi-Organ-Chip materials	29
3.1.9 Instruments	30
3.1.10 Software.....	31
3.2 Methods	32
3.2.1 Cell culture	32
3.2.2 Isolation and expansion of MSCs.....	32
3.2.3 Passaging cells.....	32
3.2.4 Isolation of HSPCs	33
3.2.5 Bone marrow model	33

3.2.6 Microfluidic system.....	36
3.2.7 Immunofluorescence and 2-photon microscopy	38
3.2.8 Scanning electron microscopy (SEM).....	38
3.2.9 Flow cytometry.....	39
3.2.10 CFU-GEMM assay	39
3.2.11 Quantitative PCR.....	40
3.2.12 In vivo experiments	40
3.2.13 Statistical analysis	41
4. Results.....	42
4.1 Building up a suitable environment.....	42
4.2 Maintenance of the niche in serum-free medium	45
4.3 HSCs maintain their native phenotype after four weeks of culture.....	52
4.4 HSPCs differentiate into various lineages	56
4.5 The bone marrow model is robust	61
4.6 MSC HSPC interaction visualized by scanning electron microscopy	63
4.7 Culture of the bone marrow model on the MOC for eight weeks	65
4.8 HSCs cultured in the bone marrow model engraft in irradiated immunocompromised mice	69
5. Discussion	76
5.1 MSCs generate an environment similar to the in vivo bone marrow niche	77
5.2 Gene expression in the bone marrow model remains stable in a serum-free environment.....	79
5.3 MSCs alter the environment of the scaffold.....	80
5.4 HSCs retain their native phenotype for four weeks.....	81
5.5 HSCs retain their native phenotype up to eight weeks.....	83
5.6 Leaving the environment of the ceramic promotes differentiation	84
5.7 Erythropoiesis and granulopoiesis is observable within the MOC	85
5.8 The model is robust	88
5.9 HSCs cultured in the bone marrow model on the MOC are capable of engrafting in irradiated immunocompromised mice	89
6. Conclusion	92
7. Perspectives	95
8. References	98
9. List of figures	111
10. List of tables	113
11. Graphical Abstract.....	114
12. Acknowledgment	115
Eidesstattliche Erklärung	116